

Vita S Forte™

Vita S Forte™ Nutritional Supplement

Each Vita S Forte caplet contains:

Fat-Soluble Vitamins

Vitamin A (acetate).....	3000 IU
Vitamin A (beta carotene).....	1000 IU
Vitamin E (dl-alpha tocopheryl acetate).....	60 IU
Vitamin D3 (ergocalciferol).....	400 IU

Water-Soluble Vitamins

Folic Acid.....	1.0 mg
Vitamin C (ascorbic acid).....	500 mg
Vitamin B1 (thiamine mononitrate).....	20 mg
Vitamin B2 (riboflavin).....	20 mg
Vitamin B6 (pyridoxine hydrochloride).....	25 mg
Vitamin B12 (cyanocobalamin).....	50 mcg
Niacin (as niacinamide).....	100 mg
Biotin.....	0.15 mg
Pantothenic Acid (calcium pantothenate).....	25 mg

Minerals

Iron (ferrous fumarate).....	10 mg
Selenium (sodium selenate).....	50 mcg
Magnesium (magnesium oxide).....	50 mg
Zinc (zinc oxide).....	15 mg
Molybdenum (sodium molybdate).....	20 mcg
Copper (cupric oxide).....	3 mg
Chromium (chromium nitrate).....	0.05 mg

Description: Description: Vita S Forte is specially formulated as a therapeutic or prophylactic nutritional supplement in stressful conditions. Vita S Forte is a multivitamin/multimineral caplet and requires a prescription.

Vita S Forte supplies therapeutic levels of water-soluble vitamins (ascorbic acid and all B complex vitamins except biotin); supplemental levels of biotin, fat-soluble vitamins (A, D and E) and minerals (iron, chromium, magnesium, copper, selenium and zinc).

Recent medical reports have shown a causal effect regarding the use of Folic Acid and the lowering of elevated levels of homocysteine. Elevated homocysteine levels have been identified as one of the risk factors for heart attacks and strokes in men.

INDICATIONS AND USAGE: Vita S Forte is indicated for therapeutic conditions or for prophylactic purposes, which include:

- stressful conditions
- conditions which cause reduced absorption or bioavailability of essential vitamins and minerals, or which cause depletion of such vitamins and minerals.
- certain conditions resulting from severe B-Vitamin or Vitamin C deficiency.
- gastrointestinal disorders, chronic or acute infections, especially those associated with febrile illness, prolonged or wasting disease, hyperthyroidism, congestive heart disease, poorly controlled diabetes, chronic alcoholism, or other physiologically stressful conditions.

(See reverse)

Vita S Forte™

- inadequate intake of vitamins and minerals due to unbalanced or highly restrictive diets such as those which can be associated with, but not limited to, any states of severe malnutrition such as anorexia.
- conditions which may benefit from increased intake of essential vitamins and minerals.
- conditions resulting from severe deficiency of Vitamin C and B-vitamins.
- patients on estrogen replacement therapy, or oral contraceptives.
- patients on a course of antibacterial drug therapy, or other drug therapy, which may affect gastrointestinal absorption of essential vitamins and minerals.
- water soluble vitamin deficiency conditions.
- patients recovering from surgery or trauma involving fractures, severe burns, or extensive tissue damage injuries.
- pregnant women, and women with heavy menstrual bleeding.

CONTRAINDICATIONS: Vita S Forte is contraindicated in patients hypersensitive to any of its components.

WARNINGS:

Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

Not intended for treatment of pernicious anemia or other megaloblastic anemias where vitamin B12 is deficient. Neurologic involvement may develop or progress, despite temporary remission of anemia, in patients with vitamin B12 deficiency who receive supplemental folic acid and who are inadequately treated with B12.

Drug and Treatment Interactions: As little as 5 mg pyridoxine daily can decrease the efficacy of levodopa in the treatment of parkinsonism. Therefore, Vita S Forte is not recommended for patients undergoing such therapy.

ADVERSE REACTIONS: Adverse reactions have been reported with specific vitamins and minerals, but generally at levels substantially higher than those in Vita S Forte. However, allergic and idiosyncratic reactions are possible at lower levels. Iron, even at the usual recommended levels, has been associated with gastrointestinal intolerance in some patients.

DOSAGE AND ADMINISTRATION: Usual adult dosage: one caplet daily.

HOW SUPPLIED: Dark green, oblong, bisected caplets, embossed 'RIS 102'

NDC #64980-102-01 Bottles of 100 caplets.

Store at controlled room temperature 15°-30°C (59°-86°F). Avoid excessive heat. Protect from light and moisture.

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.

Rx Only

Manufactured for:
Rising Pharmaceuticals, Inc.
Paramus, NJ 07652

Manufactured by:
Phoenix Labs, Inc.
Hicksville, NY 11801

Revised July 2001