

**MEDICATION GUIDE**  
**Gabapentin (GA ba PEN tin)**  
**Tablets**

Dispense with Medication Guide available at: <https://www.risingpharma.com/Medguides/GabapentinTabletsUSPMG.pdf>

**What is the most important information I should know about Gabapentin tablets?**

**Do not stop taking Gabapentin tablets without first talking to your healthcare provider.**

Stopping Gabapentin tablets suddenly can cause serious problems.

**Gabapentin tablets can cause serious side effects including:**

**1. Suicidal Thoughts.** Like other antiepileptic drugs, Gabapentin tablets may cause suicidal thoughts or actions in a very small number of people, about 1 in 500.

**Call a healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you:**

- thoughts about suicide or dying
- attempts to commit suicide
- new or worse depression
- new or worse anxiety
- feeling agitated or restless
- panic attacks
- trouble sleeping (insomnia)
- new or worse irritability
- acting aggressive, being angry, or violent
- acting on dangerous impulses
- an extreme increase in activity and talking (mania)
- other unusual changes in behavior or mood

**How can I watch for early symptoms of suicidal thoughts and actions?**

- Pay attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings.
  - Keep all follow-up visits with your healthcare provider as scheduled.

Call your healthcare provider between visits as needed, especially if you are worried about symptoms.

**Do not stop taking Gabapentin tablets without first talking to a healthcare provider.**

- Stopping Gabapentin tablets suddenly can cause serious problems. Stopping a seizure medicine suddenly in a patient who has epilepsy can cause seizures that will not stop (status epilepticus).
- Suicidal thoughts or actions can be caused by things other than medicines. If you have suicidal thoughts or actions, your healthcare provider may check for other causes.

**2. Changes in behavior and thinking** -Using Gabapentin tablets in children 3 to 12 years of age can cause emotional changes, aggressive behavior, problems with concentration, restlessness, changes in school performance, and hyperactivity.

**3. Gabapentin tablets may cause serious or life-threatening allergic reactions** that may affect your skin or other parts of your body such as your liver or blood cells. This may cause you to be hospitalized or to stop Gabapentin tablets. You may or may not have a rash with an allergic reaction caused by Gabapentin tablets. Call a healthcare provider right away if you have any of the following symptoms:

- skin rash
- hives
- difficulty breathing
- fever
- swollen glands that do not go away
- swelling of your face, lips, throat, or tongue
- yellowing of your skin or of the whites of the eyes
- unusual bruising or bleeding
- severe fatigue or weakness
- unexpected muscle pain
- frequent infections

These symptoms may be the first signs of a serious reaction. A healthcare provider should examine you to decide if you should continue taking Gabapentin tablets.

**4. Serious breathing problems.** Serious breathing problems can occur when Gabapentin tablets are taken with other medicines that can cause severe sleepiness or decreased awareness, or when it is taken by someone who already has breathing problems. Watch for increased sleepiness or decreased breathing when starting Gabapentin tablets or when the dose is increased. Get help right away if breathing problems occur.

**What are Gabapentin tablets?**

Gabapentin tablets are a prescription medicine used to treat:

- Pain from damaged nerves (postherpetic pain) that follows healing of shingles (a painful rash that comes after a herpes zoster infection) in adults.
- Partial seizures when taken together with other medicines in adults and children 3 years of age and older with seizures.

**Who should not take Gabapentin tablets?**

Do not take Gabapentin tablets if you are allergic to gabapentin or any of the other ingredients in Gabapentin tablets. See the end of this Medication Guide for a complete list of ingredients in Gabapentin tablets.

**What should I tell my healthcare provider before taking Gabapentin tablets?**

**Before taking Gabapentin tablets, tell your healthcare provider if you:**

- have or have had kidney problems or are on hemodialysis
- have or have had depression, mood problems, or suicidal thoughts or behavior
- have diabetes
- have breathing problems
- are pregnant or plan to become pregnant. It is not known if Gabapentin tablets can harm your unborn baby. Tell your healthcare provider right away if you become pregnant while taking Gabapentin tablets. You and your healthcare provider will decide if you should take Gabapentin tablets while you are pregnant.
- are breast-feeding or plan to breast-feed. Gabapentin tablets can pass into breast milk. You and your healthcare provider should decide how you will feed your baby while you take Gabapentin tablets.

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter

Size: 100 x 230 mm  
Clour :01, Black

medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take any opioid pain medicine (such as oxycodone), any medicines for anxiety (such as lorazepam) or insomnia (such as zolpidem), or any medicines that make you sleepy.

You may have a higher chance for dizziness, sleepiness, or breathing problems if these medicines are taken with Gabapentin tablets.

Taking Gabapentin tablets with certain other medicines can cause side effects or affect how well they work. Do not start or stop other medicines without talking to your healthcare provider.

Know the medicines you take. Keep a list of them and show it to your healthcare provider and pharmacist when you get a new medicine.

#### How should I take Gabapentin tablets?

- Take Gabapentin tablets exactly as prescribed. Your healthcare provider will tell you how much Gabapentin tablets to take.
  - Do not change your dose of Gabapentin tablets without talking to your healthcare provider.
  - If you take Gabapentin tablets and break a tablet in half, the unused half of the tablet should be taken at your next scheduled dose. Half tablets not used within 28 days of breaking should be thrown away.
- Gabapentin tablets can be taken with or without food. If you take an antacid containing aluminum and magnesium, such as Maalox<sup>®</sup>, Mylanta<sup>®</sup>, Gelusil<sup>®</sup>, Gaviscon<sup>®</sup>, or Di-Gel<sup>®</sup>, you should wait at least 2 hours before taking your next dose of Gabapentin tablets.

If you take too much Gabapentin tablets, call your healthcare provider or your local Poison Control Center right away at 1-800-222-1222.

#### What should I avoid while taking Gabapentin tablets?

- Do not drink alcohol or take other medicines that make you sleepy or dizzy while taking Gabapentin tablets without first talking with your healthcare provider. Taking Gabapentin tablets with alcohol or drugs that cause sleepiness or dizziness may make your sleepiness or dizziness worse.
- Do not drive, operate heavy machinery, or do other dangerous activities until you know how Gabapentin tablets affects you. Gabapentin tablets can slow your thinking and motor skills.

#### What are the possible side effects of Gabapentin tablets?

##### Gabapentin tablets may cause serious side effects including:

See "What is the most important information I should know about Gabapentin tablets?"

- problems driving while using Gabapentin tablets. See "What I should avoid while taking Gabapentin tablets?"
- sleepiness and dizziness, which could increase the occurrence of accidental injury, including falls

##### The most common side effects of Gabapentin tablets include:

- lack of coordination
- feeling drowsy
- viral infection
- nausea and vomiting
- difficulty with speaking
- jerky movements
- tremor
- difficulty with coordination
- swelling, usually of legs and feet
- double vision
- feeling tired
- unusual eye movement
- fever

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of Gabapentin tablets. For more information, ask your healthcare provider or pharmacist.

**Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**

#### How should I store Gabapentin tablets?

- Store at 25°C (77°F); excursions permitted between 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].
- Keep Gabapentin tablets and all medicines out of the reach of children.

#### General information about the safe and effective use of Gabapentin tablets

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use Gabapentin tablets for a condition for which it was not prescribed. Do not give Gabapentin tablets to other people, even if they have the same symptoms that you have. It may harm them.

This Medication Guide summarizes the most important information about Gabapentin tablets. If you would like more information, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about Gabapentin tablets that was written for healthcare professionals.

For more information, call Rising Pharma Holdings Inc, at 1-844-874-7464.

#### What are the ingredients in Gabapentin tablets?

**Active ingredient:** Gabapentin, USP

**Inactive ingredients:** Corn starch, copovidone, poloxamer, and magnesium stearate. The film coating Opadry White contains hydroxypropyl cellulose and talc.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

The brand names listed are trademarks of their respective owners and are not trademarks of the Graviti Pharmaceuticals Private Limited.

#### Manufactured by:

Graviti Pharmaceuticals Pvt Ltd.  
Telangana-502307, INDIA.

#### Manufactured for:

Rising Pharma Holdings, Inc.  
East Brunswick, NJ 08816

Mfg. Lic. No.: 12/SRD/TS/2017/F/G

Revised: 03/2024

MGR11750-00