


<div>MEDICATION GUIDE</div> <div>Mirtazapine Orally Disintegrating Tablets, USP</div> <div>(mir taz’ a peen)</div> <div>for oral use</div>				
<p><b>What is the most important information I should know about mirtazapine orally disintegrating tablets?</b></p> <p><b>Mirtazapine orally disintegrating tablets may cause serious side effects, including:</b></p> <ul style="list-style-type: none"><li>• <b>Increased risk of suicidal thoughts or actions in some children and young adults.</b> Mirtazapine orally disintegrating tablets, and other antidepressant medicines may increase suicidal thoughts or actions in some people 24 years of age and younger, <b>especially within the first few months of treatment or when the dose is changed. Mirtazapine orally disintegrating tablets are not for use in children.</b><ul style="list-style-type: none"><li>◦ Depression or other serious mental illnesses are the most important causes of suicidal thoughts or actions.</li></ul></li></ul> <p><b>How can I watch for and try to prevent suicidal thoughts and actions?</b></p> <ul style="list-style-type: none"><li>◦ Pay close attention to any changes, especially sudden changes in mood, behavior, thoughts, or feelings, or if you develop suicidal thoughts or actions. This is very important when an antidepressant medicine is started or when the dose is changed.</li><li>◦ Call your healthcare provider right away to report new or sudden changes in mood, behavior, thoughts, or feelings.</li><li>◦ Keep all follow-up visits with your healthcare provider as scheduled. Call your healthcare provider between visits as needed, especially if you have concerns about symptoms.</li></ul> <p><b>Call your healthcare provider or get emergency medical help right away if you or your family member have any of the following symptoms, especially if they are new, worse, or worry you:</b></p> <div><div><ul style="list-style-type: none"><li>• attempts to commit suicide</li><li>• acting aggressive, being angry or violent</li><li>• new or worse depression</li><li>• panic attacks</li><li>• new or worse irritability</li><li>• an extreme increase in activity or talking (mania)</li></ul></div><div><ul style="list-style-type: none"><li>• acting on dangerous impulses</li><li>• thoughts about suicide or dying</li><li>• new or worse anxiety</li><li>• feeling very agitated or restless</li><li>• trouble sleeping</li><li>• other unusual changes in behavior or mood</li></ul></div></div>				
<p><b>What are mirtazapine orally disintegrating tablets?</b></p> <p>Mirtazapine orally disintegrating tablets are prescription medicines used to treat a certain type of depression called Major Depressive Disorder (MDD) in adults. It is not known if mirtazapine orally disintegrating tablets are safe and effective for use to treat MDD in children.</p>				
<p><b>Who should not take mirtazapine orally disintegrating tablets?</b></p> <p><b>Do not take mirtazapine orally disintegrating tablets if you:</b></p> <ul style="list-style-type: none"><li>• take a Monoamine Oxidase Inhibitor (MAOI)</li><li>• have stopped taking an MAOI in the last 14 days</li><li>• are being treated with the antibiotic linezolid or intravenous methylene blue</li><li>• if you are allergic to mirtazapine or any of the ingredients in mirtazapine orally disintegrating tablets. See the end of this Medication Guide for a complete list of ingredients in mirtazapine orally disintegrating tablets.</li></ul> <p>Ask your healthcare provider or pharmacist if you are not sure if you take an MAOI, including the antibiotic linezolid or intravenous methylene blue.</p> <p><b>Do not start taking an MAOI for at least 14 days after you stop treatment with mirtazapine orally disintegrating tablets.</b></p>				
<p><b>Before taking mirtazapine orally disintegrating tablets, tell your healthcare provider about all your medical conditions, including if you:</b></p> <ul style="list-style-type: none"><li>• have a history of suicide or depression</li><li>• have a history or family history of bipolar disorder, mania or hypomania</li><li>• have a low white blood cell count</li><li>• have glaucoma (high pressure in the eye)</li><li>• have or had heart problems or stroke</li><li>• have an abnormal heart beat called QT prolongation or a family history of QT prolongation</li><li>• have seizures</li><li>• have high cholesterol or triglyceride levels</li><li>• have low sodium levels in your blood</li><li>• have or had kidney or liver problems</li><li>• have low blood pressure</li><li>• have phenylketonuria (PKU). Mirtazapine orally disintegrating tablets contain phenylalanine which is a part of aspartame.</li><li>• are pregnant or plan to become pregnant. It is not known if mirtazapine orally disintegrating tablets will harm your unborn baby.<ul style="list-style-type: none"><li>◦ Talk to your healthcare provider if you become pregnant or think you may be pregnant during treatment with mirtazapine orally disintegrating tablets.</li><li>◦ If you become pregnant while taking mirtazapine orally disintegrating tablets, talk to your healthcare provider about registering with the National Pregnancy Registry for Antidepressants. You can register by calling 1-844-405-6185 or visiting online at <a href="https://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/antidepressants/">https://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/antidepressants/</a>. The purpose of this registry is to monitor the pregnancy outcomes in women who have been treated with mirtazapine orally disintegrating tablets at any time during pregnancy.</li></ul></li><li>• are breastfeeding or plan to breastfeed. Mirtazapine may pass into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with mirtazapine orally disintegrating tablets.</li></ul> <p><b>Tell your healthcare provider about all the medicines you take,</b> including prescription and over-the-counter medicines, vitamins, and herbal supplements.</p> <p>Mirtazapine orally disintegrating tablets and other medicines may affect each other causing possible serious side effects. Mirtazapine orally disintegrating tablets may affect the way other medicines work and other medicines may affect the way mirtazapine orally disintegrating tablets work.</p> <p><b>Especially tell your healthcare provider if you take:</b></p> <ul style="list-style-type: none"><li>• MAOIs</li><li>• medicines to treat migraine headaches known as triptans</li><li>• tricyclic antidepressants</li><li>• fentanyl</li><li>• lithium</li><li>• tramadol</li><li>• tryptophan</li><li>• buspirone</li><li>• amphetamines</li><li>• benzodiazepines</li><li>• St. John’s Wort</li><li>• medicines used to treat mood, anxiety, psychotic or thought disorders, including selective serotonin reuptake inhibitors (SSRIs) and serotonin norepinephrine reuptake inhibitors (SNRIs)</li><li>• medicines that may affect your heart rhythm (such as certain antibiotics and some antipsychotics)</li></ul> <p>Ask your healthcare provider if you are not sure if you are taking any of these medicines. Your healthcare provider can tell you if it is safe to take mirtazapine orally disintegrating tablets with your other medicines.</p> <p>Do not start or stop any other medicines during treatment with mirtazapine orally disintegrating tablets without talking to your healthcare provider first. Stopping mirtazapine orally disintegrating tablets suddenly may cause you to have serious side effects. See, <b>“What are the possible side effects of mirtazapine orally disintegrating tablets?”</b></p> <p>Know the medicines you take. Keep a list of them to show to your healthcare provider and pharmacist when you get a new medicine.</p>				
<p><b>How should I take mirtazapine orally disintegrating tablets?</b></p> <ul style="list-style-type: none"><li>• Take mirtazapine orally disintegrating tablets exactly as your healthcare provider tells you to. Do not change your dose or stop taking mirtazapine orally disintegrating tablets without first talking to your healthcare provider.</li><li>• Your healthcare provider may need to change the dose of mirtazapine orally disintegrating tablets until it is the right dose for you.</li><li>• Take mirtazapine orally disintegrating tablets 1 time each day, preferably in the evening at bedtime.</li><li>• If you take too much mirtazapine call your healthcare provider or poison control center at 1-800-222-1222 right away or go to the nearest hospital emergency room.</li></ul> <p><b>How to take mirtazapine orally disintegrating tablets:</b></p> <ul style="list-style-type: none"><li>◦ Keep mirtazapine orally disintegrating tablets in the blister pack or container pack until you are ready to take them. Mirtazapine orally disintegrating tablet should be used immediately after opening the blister pack or container pack. Do not save it for future use.</li><li>◦ Use dry hands to open the blister pack or container pack.</li><li>◦ As soon as the blister pack or container pack is opened, remove mirtazapine orally disintegrating tablet right away and place it on the tongue.</li><li>◦ Place mirtazapine orally disintegrating tablet on the tongue and allow it to melt (disintegrate). <b>Do not chew, crush, or split</b> mirtazapine orally disintegrating tablets.</li><li>◦ Mirtazapine orally disintegrating tablets will disintegrate rapidly on the tongue and can be swallowed with saliva. No water is needed for taking them.</li></ul>				
<p><b>What should I avoid while taking mirtazapine orally disintegrating tablets?</b></p> <ul style="list-style-type: none"><li>• Do not drive, operate heavy machinery, or do other dangerous activities until you know how mirtazapine orally disintegrating tablets affects you. Mirtazapine orally disintegrating tablets can cause sleepiness or may affect your ability to make decisions, think clearly, or react quickly.</li><li>• Avoid drinking alcohol during treatment with mirtazapine orally disintegrating tablets.</li><li>• Avoid taking medicines used to treat anxiety, insomnia, and seizures, called benzodiazepines, during treatment with mirtazapine orally disintegrating tablets. Ask your healthcare provider if you are not sure if you take one of these medicines.</li></ul>				

A/s: 250 x 450 mm    ■ Black

<div><div>AUROBINDO</div><div>Packaging Development</div></div>		Product Name	Component	Item Code	Date & Time
		Mirtazapine OD Tablets	MG	P1529961	04.01.2022 & 04.00 pm
		Country	Version No.	Reason Of Issue	Reviewed / Approved by
		USA_Rising Health	01	Commercial	
Team Leader	Jagan	Dimensions (mm)	Colours: 01		
Initiator	Riyaz	A/s: 250 x 450 mm			
Artist	<div>Sree Designers</div>				
Additional Information: Supersede Code: P1525712 APL Ref Code: P1529778					

### What are the possible side effects of mirtazapine orally disintegrating tablets?

**Mirtazapine orally disintegrating tablets may cause serious side effects, including:**

- See, “What is the most important information I should know about mirtazapine orally disintegrating tablets?”
- **Low white blood cell count.** Tell your healthcare provider right away if you develop any signs or symptoms of a low white blood cell count, including:
  - o fever
  - o sore throat
  - o flu-like symptoms
  - o chills
  - o mouth or nose sores
  - o infections
- **Serotonin syndrome.** A potentially life-threatening problem called serotonin syndrome can happen when you take mirtazapine orally disintegrating tablets with certain other medicines. See, “**Who should not take mirtazapine orally disintegrating tablets?**” Stop taking mirtazapine orally disintegrating tablets and call your healthcare provider or go to the nearest hospital emergency room right away if you have any of the following signs and symptoms of serotonin syndrome:
  - o agitation
  - o confusion
  - o fast heart beat
  - o dizziness
  - o flushing
  - o tremors, stiff muscles, or muscle twitching
  - o seizures
  - o seeing or hearing things that are not real (hallucinations)
  - o coma
  - o blood pressure changes
  - o sweating
  - o high body temperature (hyperthermia)
  - o loss of coordination
  - o nausea, vomiting, diarrhea
- **Eye problems (angle-closure glaucoma).** Mirtazapine orally disintegrating tablets may cause a certain type of eye problem called angle-closure glaucoma. Call your healthcare provider if you have eye pain, changes in your vision, or swelling or redness in or around the eye. Only some people are at risk for these problems. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are.
- **Heart rhythm problems.**
- **Severe skin reaction.** Mirtazapine orally disintegrating tablets may cause a severe skin reaction that may include rash, fever, swollen glands, and other organ involvement such as liver, kidney, lung and heart. The reaction may sometimes be fatal. Tell your healthcare provider right away if you experience any of these signs.
- **Increased appetite and weight gain.**
- **Sleepiness.** See, “What should I avoid while taking mirtazapine orally disintegrating tablets?”
- **Mania or hypomania (manic episodes)** in people who have a history of bipolar disorder. Symptoms may include:
  - o greatly increased energy
  - o racing thoughts
  - o unusually grand ideas
  - o talking more or faster than usual
  - o severe trouble sleeping
  - o reckless behavior
  - o excessive happiness or irritability
- **Seizures (convulsions).**
- **Increased fat levels (cholesterol and triglycerides) in your blood.**
- **Low sodium levels in your blood (hyponatremia).** Low sodium levels in your blood may be serious and may cause death. Elderly people may be at greater risk for this. Signs and Symptoms of low sodium levels in your blood may include:
  - o headache
  - o memory changes
  - o weakness and unsteadiness on your feet which can lead to falls
  - o difficulty concentrating
  - o confusion
- **In severe or more sudden cases, signs and symptoms include:**
  - o hallucinations (seeing or hearing things that are not real)
  - o seizures
  - o respiratory arrest
  - o fainting
  - o coma
  - o death
- **Changes in liver function tests.**
- **Discontinuation syndrome.** Suddenly stopping mirtazapine orally disintegrating tablets may cause you to have serious side effects. Your healthcare provider may want to decrease your dose slowly. Symptoms may include:
  - o dizziness
  - o irritability and agitation
  - o anxiety
  - o sweating
  - o seizures
  - o ringing in your ears (tinnitus)
  - o nausea and vomiting
  - o problems sleeping
  - o tiredness
  - o confusion
  - o electric shock sensation (paresthesia)
  - o shaking (tremor)
  - o headache
  - o abnormal dreams
  - o changes in your mood
  - o hypomania

### How should I store mirtazapine orally disintegrating tablets?

- Store mirtazapine orally disintegrating tablets at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F).
- Keep mirtazapine orally disintegrating tablets away from light and moisture.

**Keep mirtazapine orally disintegrating tablets, and all medicines out of the reach of children.**

**General information about the safe and effective use of mirtazapine orally disintegrating tablets.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use mirtazapine orally disintegrating tablets for a condition for which it was not prescribed. Do not give mirtazapine orally disintegrating tablets to other people, even if they have the same symptoms that you have. They may harm them. You can ask your healthcare provider or pharmacist for information about mirtazapine orally disintegrating tablets that is written for healthcare professionals.

**What are the ingredients in mirtazapine orally disintegrating tablets?**

**Active ingredient:** mirtazapine

**Inactive ingredients:** aspartame, colloidal silicon dioxide, crospovidone, magnesium stearate, mannitol, microcrystalline cellulose, natural and artificial peppermint flavor, and strawberry guarana flavor.

For more information about mirtazapine orally disintegrating tablets call Rising Health, LLC at 1-833-395-6928.

Dispense with Medication Guide available at: <http://www.risingpharma.com/med-guides.html>

**Distributed by:**

**Rising Health, LLC**  
Saddle Brook, NJ 07663

**Made in India**

Code: TS/DRUGS/19/1993

This Medication Guide has been approved by the U.S. Food and Drug Administration  
Revised: 01/2022