

<p style="text-align: center;">MEDICATION GUIDE Montelukast Sodium Tablets, USP (mon te loo'kast soe'dee um)</p> <p>What is the most important information I should know about montelukast sodium tablets?</p> <p>Serious mental health problems have happened in people taking montelukast sodium tablets or even after treatment has stopped. This can happen in people with or without a history of mental health problems. Stop taking montelukast sodium tablets and tell your healthcare provider right away if you or your child have any unusual changes in behavior or thinking, including any of these symptoms:</p> <table><tr><td>• agitation, including aggressive behavior or hostility</td><td>• hallucinations (seeing or hearing things that are not really there)</td><td>• suicidal thoughts and actions (including suicide)</td></tr><tr><td>• attention problems</td><td>• memory problems</td><td>• tremor</td></tr><tr><td>• bad or vivid dreams</td><td>• obsessive-compulsive symptoms</td><td>• trouble sleeping</td></tr><tr><td>• depression</td><td>• restlessness</td><td>• uncontrolled muscle movements</td></tr><tr><td>• disorientation (confusion)</td><td>• sleep walking</td><td></td></tr><tr><td>• feeling anxious</td><td>• stuttering</td><td></td></tr><tr><td>• irritability</td><td></td><td></td></tr></table>	• agitation, including aggressive behavior or hostility	• hallucinations (seeing or hearing things that are not really there)	• suicidal thoughts and actions (including suicide)	• attention problems	• memory problems	• tremor	• bad or vivid dreams	• obsessive-compulsive symptoms	• trouble sleeping	• depression	• restlessness	• uncontrolled muscle movements	• disorientation (confusion)	• sleep walking		• feeling anxious	• stuttering		• irritability		
• agitation, including aggressive behavior or hostility	• hallucinations (seeing or hearing things that are not really there)	• suicidal thoughts and actions (including suicide)																			
• attention problems	• memory problems	• tremor																			
• bad or vivid dreams	• obsessive-compulsive symptoms	• trouble sleeping																			
• depression	• restlessness	• uncontrolled muscle movements																			
• disorientation (confusion)	• sleep walking																				
• feeling anxious	• stuttering																				
• irritability																					
<p>What are montelukast sodium tablets?</p> <p>Montelukast sodium tablets are a prescription medicine that blocks substances in the body called leukotrienes. This may help to improve symptoms of asthma and inflammation of the lining of the nose (allergic rhinitis). Montelukast sodium tablets does not contain a steroid. Montelukast sodium tablets are used to:</p> <ol style="list-style-type: none">1. Prevent asthma attacks and for the long-term treatment of asthma in adults and adolescents age 15 years and older. Do not take montelukast sodium tablets if you need relief right away for a sudden asthma attack. If you have an asthma attack, you should follow the instructions your healthcare provider gave you for treating asthma attacks.2. Prevent exercise-induced asthma in people 15 years of age and older.3. Help control the symptoms of allergic rhinitis such as sneezing, stuffy nose, runny nose, and itching of the nose. Montelukast sodium tablets are used to treat the following in people who have already taken other medicines that did not work well enough or in people who could not tolerate other medicines:<ul style="list-style-type: none">• outdoor allergies that happen part of the year (seasonal allergic rhinitis) in adults and adolescents age 15 years and older, and• indoor allergies that happen all year (perennial allergic rhinitis) in adults and adolescents age 15 years and older.																					
<p>Do not take montelukast sodium tablets if you are allergic to any of its ingredients. See the end of this Medication Guide for a complete list of the ingredients in montelukast sodium tablets.</p>																					
<p>Before taking montelukast sodium tablets, tell your healthcare provider about all your medical conditions, including if you:</p> <ul style="list-style-type: none">• are allergic to aspirin.• have or have had mental health problems.• are pregnant or plan to become pregnant. Talk to your healthcare provider if you are pregnant or plan to become pregnant, montelukast sodium tablets may not be right for you.• are breastfeeding or plan to breastfeed. It is not known if montelukast sodium passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby while taking montelukast sodium tablets. <p>Tell your healthcare provider about all the medicines you take, including prescription and over the counter medicines, vitamins, and herbal supplements. Some medicines may affect how montelukast sodium tablets works, or montelukast sodium tablets may affect how your other medicines work.</p>																					
<p>How should I take montelukast sodium tablets?</p> <p>For anyone who takes montelukast sodium tablets:</p> <ul style="list-style-type: none">• Take montelukast sodium tablets exactly as prescribed by your healthcare provider. Your healthcare provider will tell you how much montelukast sodium tablets to take, and when to take it.• Stop taking montelukast sodium tablets and tell your healthcare provider right away if you or your child have any unusual changes in behavior or thinking.• You can take montelukast sodium tablets with food or without food.• If you or your child misses a dose of montelukast sodium tablets, just take the next dose at your regular time. Do not take 2 doses at the same time.• If you take too much montelukast sodium tablets, call your healthcare provider right away. <p>For adults and adolescents 15 years of age and older with asthma:</p> <ul style="list-style-type: none">• Take montelukast sodium tablets 1 time each day, in the evening. Continue to take montelukast sodium tablets every day for as long as your healthcare provider prescribes it, even if you have no asthma symptoms.• Tell your healthcare provider right away if your asthma symptoms get worse, or if you need to use your rescue inhaler medicine more often for asthma attacks.• Always have your rescue inhaler medicine with you for asthma attacks.• Continue to take your other asthma medicines as prescribed unless your healthcare provider tells you to change how you take these medicines. <p>For people 15 years of age and older for the prevention of exercise-induced asthma:</p> <ul style="list-style-type: none">• Take montelukast sodium tablets at least 2 hours before exercise.• Always have your rescue inhaler medicine with you for asthma attacks.																					

Size: 100 x 230 mm
Clour :01, Black

