

Dispense with Medication Guide available at: <http://www.risingpharma.com/med-guides.html>

**Medication Guide**  
**Sertraline Hydrochloride Oral Solution USP**  
 (ser' tra leen hye'' droe klor' ide)

**What is the most important information I should know about sertraline hydrochloride oral solution?**

**Sertraline hydrochloride oral solution and other antidepressant medicines may cause serious side effects. Call your healthcare provider right away if you have any of the following symptoms, or call 911 if there is an emergency.**

- Suicidal thoughts or actions:**
  - Sertraline hydrochloride oral solution and other antidepressant medicines may increase suicidal thoughts or actions** in some people 24 years of age and younger, especially within the **first few months of treatment or when the dose is changed.**
  - Depression or other serious mental illnesses are the most important causes of suicidal thoughts or actions.
  - Watch for these changes and call your healthcare provider right away if you notice new or sudden changes in mood, behavior, actions, thoughts, or feelings, especially if severe.
    - Pay particular attention to such changes when sertraline hydrochloride oral solution is started or when the dose is changed.
    - Keep all follow-up visits with your healthcare provider and call between visits if you are worried about symptoms.

**Call your healthcare provider right away if you have any of the following symptoms, or call 911 if an emergency, especially if they are new, worse, or worry you:**

- |   |   |
|---|---|
| o attempts to commit suicide  | o acting on dangerous impulses              |
| o acting aggressive or violent  | o thoughts about suicide or dying           |
| o new or worse depression   | o new or worse anxiety or panic attacks     |
| o feeling agitated, restless, angry or irritable                      | o trouble sleeping                          |
| o an increase in activity or talking more than what is normal for you | o other unusual changes in behavior or mood |

- Serotonin Syndrome.** This condition can be life-threatening and symptoms may include:

- |  |                                 |
|--|---------------------------------|
| • agitation, hallucinations, coma, or other changes in mental status | • nausea, vomiting, or diarrhea |
| • racing heartbeat, high or low blood pressure                       | • sweating or fever             |
| • coordination problems or muscle twitching (overactive reflexes)    | • muscle rigidity               |

- Increased chance of bleeding:** sertraline hydrochloride oral solution and other antidepressant medicines may increase your risk of bleeding or bruising, especially if you take the blood thinner warfarin (Coumadin®, Jantoven®), a non-steroidal anti-inflammatory drug (NSAIDs, like ibuprofen or naproxen), or aspirin.

- Manic episodes.** Symptoms may include:

- |                            |                           |                                       |
|----------------------------|---------------------------|---------------------------------------|
| • greatly increased energy | • severe trouble sleeping | • excessive happiness or irritability |
| • racing thoughts          | • reckless behavior       | • talking more or faster than usual   |
| • unusually grand ideas    |                           |                                       |

- Seizures or convulsions.**

- Glaucoma (angle-closure glaucoma).** Many antidepressant medicines including sertraline hydrochloride oral solution may cause a certain type of eye problem called angle-closure glaucoma. Call your healthcare provider if you have eye pain, changes in your vision, or swelling or redness in or around the eye. Only some people are at risk for these problems. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are.

- Changes in appetite or weight.** Children and adolescents should have height and weight monitored during treatment.

- Low salt (sodium) levels in the blood.** Elderly people may be at greater risk for this. Symptoms may include:

- headache
- weakness or feeling unsteady
- confusion, problems concentrating or thinking, or memory problems

- Sexual problems (dysfunction).** Taking selective serotonin reuptake inhibitors (SSRIs), including sertraline hydrochloride oral solution, may cause sexual problems.

Symptoms in males may include:

- Delayed ejaculation or inability to have an ejaculation
- Decreased sex drive
- Problems getting or keeping an erection

Symptoms in females may include:

- Decreased sex drive
- Delayed orgasm or inability to have an orgasm

Talk to your healthcare provider if you develop any changes in your sexual function or if you have any questions or concerns about sexual problems during treatment with sertraline hydrochloride oral solution. There may be treatments your healthcare provider can suggest.

**Do not stop sertraline hydrochloride oral solution without first talking to your healthcare provider.** Stopping sertraline hydrochloride oral solution too quickly may cause serious symptoms including:

- anxiety, irritability, high or low mood, feeling restless or changes in sleep habits
- headache, sweating, nausea, dizziness
- electric shock-like sensations, shaking, confusion

**What is Sertraline Hydrochloride Oral Solution?**

Sertraline hydrochloride oral solution is a prescription medicine used to treat:

- |                                   |  |
|-----------------------------------|--|
| • Major Depressive Disorder (MDD) | • Obsessive Compulsive Disorder (OCD)    |
| • Panic Disorder                  | • Posttraumatic Stress Disorder (PTSD)   |
| • Social Anxiety Disorder         | • Premenstrual Dysphoric Disorder (PMDD) |

It is important to talk with your healthcare provider about the risks of treating depression and also the risks of not treating it. You should discuss all treatment choices with your healthcare provider.

Sertraline hydrochloride oral solution is safe and effective in treating children with OCD age 6 to 17 years.

It is not known if sertraline hydrochloride oral solution is safe and effective for use in children under 6 years of age with OCD or children with other behavior health conditions.

Talk to your healthcare provider if you do not think that your condition is getting better with sertraline hydrochloride oral solution treatment.

**Who should not take sertraline hydrochloride oral solution?**


**Do not take sertraline hydrochloride oral solution if you:**

- take a monoamine oxidase inhibitor (MAOI). Ask your healthcare provider or pharmacist if you are not sure if you take an MAOI, including the antibiotic linezolid.
- have taken an MAOI within 2 weeks of stopping sertraline hydrochloride oral solution unless directed to do so by your healthcare provider.
- have stopped taking an MAOI in the last 2 weeks unless directed to do so by your healthcare provider.
- take any other medicines that contain sertraline (such as sertraline HCl or sertraline hydrochloride).
- take the antipsychotic medicine pimozide (Orap®) because this can cause serious heart problems.
- are allergic to sertraline or any of the ingredients in sertraline hydrochloride oral solution. See the end of this Medication Guide for a complete list of ingredients in sertraline hydrochloride oral solution.
- take Antabuse® (disulfiram) (if you are taking the liquid form of sertraline hydrochloride oral solution) due to the alcohol content.

**People who take sertraline hydrochloride oral solution close in time to an MAOI may have serious or even life-threatening side effects. Get medical help right away if you have any of these symptoms:**

- |              |                              |                 |
|--------------|------------------------------|-----------------|
| o high fever | o uncontrolled muscle spasms | o stiff muscles |
|--------------|------------------------------|-----------------|

A/s: 200 x 340 mm ■ Black

	<b>Product Name</b>	<b>Component</b>	<b>Item Code</b>	<b>Date &amp; Time</b>
	Sertraline Hydrochloride OS USP	MG	P1535334	22.09.2023 & 12.30 pm
	<b>Country</b>	<b>Version No.</b>	<b>Reason Of Issue</b>	
	USA_Rising Pharma	00	Commercial	
<b>Team Leader</b>	Jagan	<b>Dimensions (mm)</b>	<b>Colours: 01</b>	
<b>Initiator</b>	Dastagiri	A/s: 200 x 340 mm		
<b>Artist</b>	Sree Designers			
<b>Additional Information:</b> Supersede Code: P1533569 APL REF Code: P1535203				

- o rapid changes in heart rate or blood pressure
- o confusion
- o loss of consciousness (pass out)

**What should I tell my healthcare provider before taking sertraline hydrochloride oral solution?**

**Before starting sertraline hydrochloride oral solution, tell your healthcare provider:**

- **if you have:**
  - o liver problems
  - o kidney problems
  - o a history of a stroke
  - o heart problems
  - o or have had seizures or convulsions
  - o high blood pressure
  - o bipolar disorder or mania
  - o low sodium levels in your blood
  - o or have had bleeding problems
- **are pregnant or plan to become pregnant.** Your baby may have withdrawal symptoms after birth or may be at increased risk for a serious lung problem at birth. Talk to your healthcare provider about the benefits and risks of taking sertraline hydrochloride oral solution during pregnancy.
  - o There is a pregnancy registry for females who are exposed to sertraline hydrochloride oral solution during pregnancy. The purpose of the registry is to collect information about the health of females exposed to sertraline hydrochloride oral solution and their baby. If you or your child become pregnant during treatment with sertraline hydrochloride oral solution, talk to your healthcare provider about registering with the National Pregnancy Registry for Antidepressants. You can register by calling 1-866-961-2388 or by visiting online at <https://womensmentalhealth.org/research/pregnancyregistry/antidepressants/>.
- **are breastfeeding or plan to breastfeed.** A small amount of sertraline hydrochloride oral solution may pass into your breast milk. Talk to your healthcare provider about the best way to feed your baby while taking sertraline hydrochloride oral solution.

**Tell your healthcare provider about all the medicines that you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**Especially tell your healthcare provider if you or your child take:**

- medicines used to treat migraine headaches called triptans
- tricyclic antidepressants
- lithium
- tramadol, fentanyl, meperidine, methadone, or other opioids
- tryptophan
- buspirone
- amphetamines
- phenytoin
- St. John's Wort
- medicines that can affect blood clotting such as aspirin, nonsteroidal anti-inflammatory drugs (NSAIDs), other antiplatelet medicines, warfarin, and other anticoagulants
- diuretics
- medicines used to treat mood, anxiety, psychotic, or thought disorders, including selective serotonin reuptake (SSRIs) and serotonin norepinephrine reuptake inhibitors (SNRIs)

Sertraline hydrochloride oral solution and some medicines may interact with each other, may not work as well, or may cause serious side effects.

Your healthcare provider or pharmacist can tell you if it is safe to take sertraline hydrochloride oral solution with your other medicines. **Do not** start or stop any medicine while taking sertraline hydrochloride oral solution without talking to your healthcare provider first.

**How should I take sertraline hydrochloride oral solution?**

- Take sertraline hydrochloride oral solution exactly as prescribed. Your healthcare provider may need to change the dose of sertraline hydrochloride oral solution until it is the right dose for you.
- Sertraline hydrochloride oral solution may look cloudy or hazy after mixing, this is normal.
- Sertraline hydrochloride oral solution must be diluted before use:
  - o **Do not** mix sertraline hydrochloride oral solution until you are ready to take it.
  - o When diluting sertraline hydrochloride oral solution, use **only** water, ginger ale, lemon/lime soda, lemonade, or orange juice.
- If you miss a dose of sertraline hydrochloride oral solution, take the missed dose as soon as you remember. If it is almost time for the next dose, skip the missed dose and take your next dose at the regular time. **Do not** take two doses of sertraline hydrochloride oral solution at the same time.

**If you take too much sertraline hydrochloride oral solution, call your healthcare provider or poison control center right away, or go to the nearest hospital emergency room right away.**

**What should I avoid while taking sertraline hydrochloride oral solution?**

Sertraline hydrochloride oral solution can cause sleepiness or may affect your ability to make decisions, think clearly, or react quickly. You should not drive, operate heavy machinery, or do other dangerous activities until you know how sertraline hydrochloride oral solution affects you. **Do not** drink alcohol while you take sertraline hydrochloride oral solution.

**What are the possible side effects of sertraline hydrochloride oral solution?**

Sertraline hydrochloride oral solution may cause serious side effects, including:

- **See “What is the most important information I should know about sertraline hydrochloride oral solution?”**

**The most common side effects in adults who take sertraline hydrochloride oral solution include:**

- nausea, loss of appetite, diarrhea, or indigestion
- change in sleep habits including increased sleepiness or insomnia
- increased sweating
- sexual problems including decreased libido and ejaculation failure
- tremor or shaking
- feeling tired or fatigued
- agitation
- anxiety

**The most common side effects in children and adolescents who take include** abnormal increase in muscle movement or agitation, nose bleeds, urinary incontinence, aggressive reaction, possible slowed growth rate, and weight change. Your child's height and weight should be monitored during treatment with sertraline hydrochloride oral solution.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of sertraline hydrochloride oral solution. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

**How should I store sertraline hydrochloride oral solution?**

- Store sertraline hydrochloride oral solution at room temperature, 68° to 77°F (20° to 25°C).
- Keep sertraline hydrochloride oral solution bottle closed tightly.

**Keep sertraline hydrochloride oral solution and all medicines out of the reach of children.**

**General information about the safe and effective use of sertraline hydrochloride oral solution**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use sertraline hydrochloride oral solution for a condition for which it was not prescribed. Do not give sertraline hydrochloride oral solution to other people, even if they have the same condition. It may harm them.

This Medication Guide summarizes the most important information about sertraline hydrochloride oral solution. If you would like more information, talk with your healthcare provider. You may ask your healthcare provider or pharmacist for information about sertraline hydrochloride oral solution that is written for healthcare professionals.

For more information call 1-866-562-4597.

**What are the ingredients in sertraline hydrochloride oral solution?**

**Active ingredient:** sertraline hydrochloride

**Inactive ingredients:** alcohol (12%), natural peppermint flavor, propylene glycol and purified water.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

All brands listed are the trademarks of their respective owners and are not trademarks of Aurobindo Pharma Limited.

**Distributed by:**

Rising Pharmaceuticals, Inc.  
East Brunswick, NJ 08816

**Made in India**

Code: TS/DRUGS/19/1993

Revised: 09/2023