



<p><b>How should I take topiramate tablets?</b></p> <ul style="list-style-type: none"><li>• Take topiramate tablets exactly as prescribed.</li><li>• Your healthcare provider may change your dose. <b>Do not</b> change your dose without talking to your healthcare provider.</li><li>• Take topiramate tablets whole. <b>Do not</b> chew the tablets. They may leave a bitter taste.</li><li>• Topiramate tablets can be taken before, during, or after a meal. Drink plenty of fluids during the day. This may help prevent kidney stones while taking topiramate tablets.</li><li>• If you take too much topiramate, call your healthcare provider right away or go to the nearest emergency room.</li><li>• If you miss a single dose of topiramate tablets, take it as soon as you can. However, if you are within 6 hours of taking your next scheduled dose, wait until then to take your usual dose of topiramate tablets, and skip the missed dose. <b>Do not</b> double your dose. If you have missed more than one dose, you should call your healthcare provider for advice.</li><li>• <b>Do not</b> stop taking topiramate tablets without talking to your healthcare provider. Stopping topiramate tablets suddenly may cause serious problems. If you have epilepsy and you stop taking topiramate tablets suddenly, you may have seizures that do not stop. Your healthcare provider will tell you how to stop taking topiramate tablets slowly.</li><li>• Your healthcare provider may do blood tests while you take topiramate tablets.</li></ul>
<p><b>What should I avoid while taking topiramate tablets?</b></p> <ul style="list-style-type: none"><li>• You should not drink alcohol while taking topiramate tablets. Topiramate tablets and alcohol can affect each other causing side effects such as sleepiness and dizziness.</li><li>• Do not drive a car or operate machinery until you know how topiramate tablets affect you. Topiramate tablets can slow your thinking and motor skills, and may affect vision.</li></ul>
<p><b>What are the possible side effects of topiramate tablets?</b></p> <p><b>Topiramate tablets may cause serious side effects including:</b></p> <p>See “What is the most important information I should know about topiramate tablets?”</p> <ul style="list-style-type: none"><li>• <b>High blood ammonia levels.</b> High ammonia in the blood can affect your mental activities, slow your alertness, make you feel tired, or cause vomiting. This has happened when topiramate tablets are taken with a medicine called valproic acid (DEPAKENE and DEPAKOTE).</li><li>• <b>Effects on thinking and alertness.</b> Topiramate tablets may affect how you think and cause confusion, problems with concentration, attention, memory, or speech. Topiramate tablets may cause depression or mood problems, tiredness, and sleepiness.</li><li>• <b>Dizziness or loss of muscle coordination.</b></li><li>• <b>Serious skin reactions.</b> Topiramate tablets may cause a severe rash with blisters and peeling skin, especially around the mouth, nose, eyes, and genitals (Stevens-Johnson syndrome). Topiramate tablets may also cause a rash with blisters and peeling skin over much of the body that may cause death (toxic epidermal necrolysis). Call your healthcare provider right away if you develop a skin rash or blisters.</li><li>• <b>Kidney stones.</b> Drink plenty of fluids when taking topiramate tablets to decrease your chances of getting kidney stones.</li><li>• <b>Low body temperature.</b> Taking topiramate tablets when you are also taking valproic acid can cause a drop in body temperature to less than 95°F, or can cause tiredness, confusion, or coma.</li></ul> <p>Call your healthcare provider right away if you have any of the symptoms above.</p> <p>The most common side effects of topiramate tablets include:</p> <ul style="list-style-type: none"><li>• tingling of the arms and legs (paresthesia)</li><li>• not feeling hungry</li><li>• nausea</li><li>• a change in the way foods taste</li><li>• diarrhea</li><li>• weight loss</li><li>• nervousness</li><li>• upper respiratory tract infection</li><li>• speech problems</li><li>• tiredness</li><li>• dizziness</li><li>• sleepiness/drowsiness</li><li>• slow reactions</li><li>• difficulty with memory</li><li>• pain in the abdomen</li><li>• fever</li><li>• abnormal vision</li><li>• decreased feeling or sensitivity, especially in the skin</li></ul> <p>Tell your healthcare provider about any side effect that bothers you or that does not go away. These are not all the possible side effects of topiramate tablets. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Rising Pharma Holdings, Inc. at 1(844) 874-7464.</p>
<p><b>How should I store topiramate tablets?</b></p> <ul style="list-style-type: none"><li>• Store topiramate tablets at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F).</li><li>• Keep topiramate tablets in a tightly closed container.</li><li>• Keep topiramate tablets dry and away from moisture.</li></ul> <p><b>Keep topiramate tablets and all medicines out of the reach of children.</b></p>
<p><b>General information about the safe and effective use of topiramate tablets.</b></p> <p>Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use topiramate tablets for a condition for which it was not prescribed. Do not give topiramate tablets to other people, even if they have the same symptoms that you have. They may harm them.</p> <p>You can ask your pharmacist or healthcare provider for information about topiramate tablets that is written for health professionals.</p>
<p><b>What are the ingredients in topiramate tablets?</b></p> <p><b>Active ingredient:</b> topiramate</p> <p><b>Inactive ingredients:</b> hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, pregelatinized starch, sodium starch glycolate, and titanium dioxide. In addition, the 50 mg and 100 mg tablets contain iron oxide yellow and 200 mg tablet contains iron oxide red.</p> <p>The brands listed are trademarks of their respective owners and are not trademarks of Aurobindo Pharma Limited.</p> <p>For more information, call Rising Pharma Holdings, Inc. at 1(844) 874-7464.</p> <p><b>Distributed by:</b> Rising Pharma Holdings, Inc. East Brunswick, NJ 08816</p> <p><b>Made in India</b></p> <p>Code: TS/DRUGS/19/1993</p>

This Medication Guide has been approved by the U.S. Food and Drug Administration.  
Revised: 07/2023