

**MEDICATION GUIDE**  
**Zolpidem Tartrate Tablets, USP**   
**(zol' pi dem tar' trate)**

Read the Medication Guide that comes with zolpidem tartrate before you start taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your healthcare provider about your medical condition or treatment.

**What is the most important information I should know about zolpidem tartrate?**

- **Do not take more zolpidem tartrate than prescribed.**
- **Do not take zolpidem tartrate unless you are able to stay in bed a full night (7 to 8 hours) before you must be active again.**
- **Take zolpidem tartrate right before you get in bed, not sooner.**

**Zolpidem tartrate may cause serious side effects, including:**

- **After taking zolpidem tartrate, you may get up out of bed while not being fully awake and do an activity that you do not know you are doing. The next morning, you may not remember that you did anything during the night.** You have a higher chance for doing these activities if you drink alcohol or take other medicines that make you sleepy with zolpidem tartrate. Reported activities include:
  - o driving a car (“sleep-driving”)
  - o making and eating food
  - o talking on the phone
  - o having sex
  - o sleep-walking

**Call your healthcare provider right away if you find out that you have done any of the above activities after taking zolpidem tartrate.**

**Do not take zolpidem tartrate if you:**

- drank alcohol that evening or before bed
- took another medicine to help you sleep.

**What is zolpidem tartrate?**

Zolpidem tartrate is a sedative-hypnotic (sleep) medicine. Zolpidem tartrate is used in adults for the short-term treatment of a sleep problem called insomnia (trouble falling asleep).

It is not known if zolpidem tartrate is safe and effective in children under the age of 18 years.

Zolpidem tartrate is a federally controlled substance (C-IV) because it can be abused or lead to dependence. Keep zolpidem tartrate in a safe place to prevent misuse and abuse. Selling or giving away zolpidem tartrate may harm others, and is against the law. Tell your healthcare provider if you have ever abused or have been dependent on alcohol, prescription medicines or street drugs.

**Who should not take zolpidem tartrate?**

- Do not take zolpidem tartrate if you are allergic to zolpidem or any other ingredients in zolpidem tartrate. See the end of this Medication Guide for a complete list of ingredients in zolpidem tartrate.

- Do not take zolpidem tartrate if you have had an allergic reaction to drugs containing zolpidem, such as Ambien CR, Edluar, Zolpimist, or Intermezzo.

Symptoms of a serious allergic reaction to zolpidem can include:

- swelling of your face, lips, and throat that may cause difficulty breathing or swallowing

**What should I tell my healthcare provider before taking zolpidem tartrate?**

**Zolpidem tartrate may not be right for you. Before starting zolpidem tartrate, tell your healthcare provider about all of your health conditions, including if you:**

- have a history of depression, mental illness, or suicidal thoughts
- have a history of drug or alcohol abuse or addiction
- have kidney or liver disease
- have a lung disease or breathing problems
- are pregnant, planning to become pregnant. It is not known if zolpidem tartrate will harm your unborn baby.
- are breastfeeding or plan to breastfeed. Zolpidem tartrate can pass into your breast milk. It is not known if zolpidem tartrate will harm your baby. Talk to your healthcare provider about the best way to feed your baby while you take zolpidem tartrate.

**Tell your healthcare provider about all of the medicines you take, including prescription and nonprescription medicines, vitamins and herbal supplements.**



Medicines can interact with each other, sometimes causing serious side effects. **Do not take zolpidem tartrate with other medicines that can make you sleepy unless your healthcare provider tells you to.**

Know the medicines you take. Keep a list of your medicines with you to show your healthcare provider and pharmacist each time you get a new medicine.

**How should I take zolpidem tartrate?**

- See **“What is the most important information I should know about zolpidem tartrate?”**
- Take zolpidem tartrate exactly as prescribed. Only take 1 zolpidem tartrate tablet a night if needed.
- Do not take zolpidem tartrate if you drank alcohol that evening or before bed.
- You should not take zolpidem tartrate with or right after a meal. Zolpidem tartrate may help you fall asleep faster if you take it on an empty stomach.

A/s: 160 x 240 mm ■ Black

		Product Name	Component	Item Code	Date & Time
		Zolpidem Tartrate Tablets	Medication Guide	P1514851	13.11.2017 & 10:00 AM
Team Leader: Ramesh Sajja Initiator: Ayyappa		Customer / Country	Version No.	Reason Of Issue	Reviewed / Approved by          Sign / Date
		Rising_Pharma_USA	04	NEW	
Artist: 		Dimensions	No. of Colours : 01		
		160 x 240 mm			
Additional Information :		Pharmacode	NA		

- Call your healthcare provider if your insomnia worsens or is not better within 7 to 10 days. This may mean that there is another condition causing your sleep problem.
- If you take too much zolpidem tartrate or overdose, get emergency treatment.

**What are the possible side effects of zolpidem tartrate?**

**Zolpidem tartrate may cause serious side effects, including:**

- **getting out of bed while not being fully awake and do an activity that you do not know you are doing.** See “**What is the most important information I should know about zolpidem tartrate?**”
- **abnormal thoughts and behavior.** Symptoms include more outgoing or aggressive behavior than normal, confusion, agitation, hallucinations, worsening of depression, and suicidal thoughts or actions.
- **memory loss**
- **anxiety**
- **severe allergic reactions.** Symptoms include swelling of the tongue or throat, and trouble breathing. Get emergency medical help if you get these symptoms after taking zolpidem tartrate.
- **falls, which may lead to severe injuries**

**Call your healthcare provider right away if you have any of the above side effects or any other side effects that worry you while using zolpidem tartrate.**

**The most common side effects of zolpidem tartrate are:**

- drowsiness
- dizziness
- diarrhea
- grogginess or feeling as if you have been drugged

**After you stop taking a sleep medicine,** you may have symptoms for 1 to 2 days such as:

- trouble sleeping
- nausea
- flushing
- lightheadedness
- uncontrolled crying
- vomiting
- stomach cramps
- panic attack
- nervousness
- stomach area pain

These are not all the side effects of zolpidem tartrate. Ask your healthcare provider or pharmacist for more information.

Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store zolpidem tartrate?**

- Store zolpidem tartrate at room temperature 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F).

**Keep zolpidem tartrate and all medicines out of reach of children.**

**General Information about the safe and effective use of zolpidem tartrate**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use zolpidem tartrate for a condition for which it was not prescribed. Do not share zolpidem tartrate with other people, even if they have the same symptoms that you have. It may harm them and it is against the law.

This Medication Guide summarizes the most important information about zolpidem tartrate. If you would like more information, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about zolpidem tartrate that is written for healthcare professionals.

For more information, call Rising Pharmaceuticals, Inc. at 1-866-562-4597.

**What are the ingredients in zolpidem tartrate tablets?**

**Active Ingredient:** Zolpidem tartrate

**Inactive Ingredients:** lactose monohydrate, microcrystalline cellulose, sodium starch glycolate, magnesium stearate, hypromellose, polyethylene glycol, and titanium dioxide.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

All brands listed are the trademarks of their respective owners and are not trademarks of Aurobindo Pharma Limited.

Dispense with Medication Guide available at: <http://www.risingpharma.com/med-guides.html>

**Distributed by:**

Rising Pharmaceuticals, Inc.  
Saddle Brook, NJ 07663

**Made in India**

Code: TS/DRUGS/22/2009

Issued: 11/2017