

Dispense with Medication Guide available at:
www.risingpharma.com/Medguides/diclofenac-sodium-delayed-release-tablets-bluepoint.pdf

**Diclofenac Sodium (dye kloe' fen ak soe' dee um)
Delayed-release Tablets, USP
Medication Guide for Nonsteroidal
Anti-inflammatory Drugs (NSAIDs)**

**What is the most important information I
should know about medicines called
Nonsteroidal Anti-inflammatory Drugs
(NSAIDs)?**

**NSAIDs can cause serious side effects,
including:**

- **Increased risk of a heart attack or stroke that
can lead to death.**

This risk may happen early in treatment and may
increase:

- with increasing doses of NSAIDs
- with longer use of NSAIDs

**Do not take NSAIDs right before or after a heart
surgery called a "coronary artery bypass graft
(CABG)."**

**Avoid taking NSAIDs after a recent heart
attack, unless your healthcare provider tells
you to. You may have an increased risk of
another heart attack if you take NSAIDs after a
recent heart attack.**

- **Increased risk of bleeding, ulcers, and tears**

**(perforation) of the esophagus (tube leading
from the mouth to the stomach), stomach and
intestines:**

- anytime during use
- without warning symptoms
- that may cause death

**The risk of getting an ulcer or bleeding
increases with:**

- past history of stomach ulcers, or stomach or
intestinal bleeding with use of NSAIDs
- taking medicines called "corticosteroids",
"anticoagulants", "SSRIs", or "SNRIs"
- increasing doses of NSAIDs
- longer use of NSAIDs
- smoking
- drinking alcohol
- older age
- poor health
- advanced liver disease
- bleeding problems

NSAIDs should only be used:

- exactly as prescribed
- at the lowest dose possible for your treatment
- for the shortest time needed

What are NSAIDs?

NSAIDs are used to treat pain and redness,
swelling, and heat (inflammation) from medical
conditions such as different types of arthritis,
menstrual cramps, and other types of short-term

pain.

Who should not take NSAIDs?

Do not take NSAIDs:

- if you had an asthma attack, hives, or
other allergic reaction with aspirin or any other
NSAIDs.
- right before or after heart bypass surgery.

**Before taking NSAIDs, tell your healthcare
provider about all of your medical conditions,
including if you:**

- have liver or kidney problems
- have high blood pressure
- have asthma
- are pregnant or plan to become pregnant.
Taking NSAIDs at about 20 weeks of
pregnancy or later may harm your unborn baby.
If you need to take NSAIDs for more than 2
days when you are between 20 and 30 weeks
of pregnancy, your healthcare provider may
need to monitor the amount of fluid in your
womb around your baby. **You should not take
NSAIDs after about 30 weeks of pregnancy.**

- are breastfeeding or plan to breastfeed.

**Tell your healthcare provider about all of the
medicines you take, including prescription or
over-the-counter medicines, vitamins or
herbal supplements.** NSAIDs and some other
medicines can interact with each other and cause
serious side effects. **Do not start taking any new
medicine without talking to your healthcare**

250 mm

120 mm

provider first.

What are the possible side effects of NSAIDs? NSAIDs can cause serious side effects, including:

See “What is the most important information I should know about medicines called Nonsteroidal Anti-inflammatory Drugs (NSAIDs)?”

- new or worse high blood pressure
- heart failure
- liver problems including liver failure
- kidney problems including kidney failure
- low red blood cells (anemia)
- life-threatening skin reactions
- life-threatening allergic reactions

• Other side effects of NSAIDs include: stomach pain, constipation, diarrhea, gas, heartburn, nausea, vomiting, and dizziness.

Get emergency help right away if you have any of the following symptoms:

- shortness of breath or trouble breathing
- chest pain
- weakness in one part or side of your body
- slurred speech
- swelling of the face or throat

Stop taking your NSAID and call your healthcare provider right away if you get any of the following symptoms:

- nausea
- more tired or weaker than usual
- diarrhea

- itching
- your skin or eyes look yellow
- indigestion or stomach pain
- flu-like symptoms
- vomit blood
- there is blood in your bowel movement or it is black and sticky like tar
- unusual weight gain
- skin reaction or blisters with fever
- swelling of the arms, legs, hands and feet

If you take too much of your NSAID, call your healthcare provider or get medical help right away.

These are not all the possible side effects of NSAIDs. For more information, ask your healthcare provider or pharmacist about NSAIDs. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Other information about NSAIDs

- Aspirin is an NSAID medicine but it does not increase the chance of a heart attack. Aspirin can cause bleeding in the brain, stomach, and intestines. Aspirin can also cause ulcers in the stomach and intestines.
- Some NSAIDs are sold in lower doses without a prescription (over-the-counter). Talk to your healthcare provider before using over-the-counter NSAIDs for more than 10 days.

General information about the safe and effective use of NSAIDs

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use NSAIDs for a condition for which it was not prescribed. Do not give NSAIDs to other people, even if they have the same symptoms that you have. It may harm them.

If you would like more information about NSAIDs, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about NSAIDs that is written for health professionals.

**Manufactured by:
UNIQUE PHARMACEUTICAL
LABORATORIES**

(A Div. of J. B. Chemicals & Pharmaceuticals Ltd.)
Mumbai 400 030, India

For BluePoint Laboratories



For more information, contact 1-844-874-7464

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Rev. 02/25

141092

250 mm

120 mm